



O-74

## EVALUATION OF AN ATTRACT-AND-KILL STRATEGY FOR THE MANAGEMENT OF THE BROWN MARMORATED STINK BUG IN NORTHERN ITALY

Antonio Masetti<sup>1</sup>, Michele Preti<sup>2</sup>, Marco Albertini<sup>3</sup>, Michele Capriotti<sup>4</sup>, Olmo Scagliarini<sup>5</sup>, Maria Grazia Tommasini<sup>6</sup>

<sup>1</sup> Department of Agricultural and Food Science – University of Bologna, v.le G. Fanin, 42 - 40127 Bologna, Italy. e-mail: antonio.masetti@unibo.it

<sup>2</sup> ASTRA Innovazione e Sviluppo, v. Tebano, 45 - 48018 Faenza (RA), Italy

<sup>3</sup> AgriTeS, v. G. Marconi, 4/2 - 40057 Granarolo dell'Emilia (BO), Italy

<sup>4</sup> Terremerse, v. Cà del Vento, 21 - 48012 Bagnacavallo (RA), Italy

<sup>5</sup> Centro Agricoltura e Ambiente "Giorgio Nicoli", v. Sant'Agata 835 - 40014 Crevalcore (BO), Italy

<sup>6</sup> RI.NOVA, v. dell'Arrigoni, 120 - 47522 Cesena (FC), Italy

**Abstract:** Attract-and-kill (AK) strategies rely on the combination of an attractant and an insecticide or a killing method to suppress pest insects. In comparison with full block sprays with broad-spectrum insecticides, AK minimize the contact between pesticides and crops resulting in many benefits from the standpoints of safety for farmers and consumers, reduction of residues on agricultural products and overall impact on the environment.

The brown marmorated stink bug (BMSB), *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae), is an Asian insect pest, extremely invasive and polyphagous, which has become very common in the USA and Southern Europe. In most countries where it has been accidentally introduced, BMSB is now a major threat to agriculture causing severe economic losses especially in tree fruit crops. Current management strategies, which mostly rely on neurotoxic broad-spectrum insecticides, are neither fully effective nor environmentally sustainable.

The implementation of AK techniques for BMSB management has been investigated in the USA resulting in acceptable reduction of crop damage, although economic issues were pointed out. Because of the severe impact of BMSB on orchards in Northern Italy, in 2021 the effectiveness of AK techniques against BMSB was evaluated in intensive friculture areas of Emilia-Romagna region. A wide area approach was used to tackle the dispersal capacity and broad range of host plants of BMSB nymphs and adults. Given that AK cannot be considered a stand-alone method, standard IPM spray programs recommended in Northern Italy for BMSB management were also carried out in the whole study areas, including controls.

Trinet® pyramid-like structures (BASF Inc.), which are made by an aluminium stand covered with a long-lasting insecticide net (LLIN) containing alpha-cypermethrin at 0.1 g m<sup>-2</sup>, were used as AK stations (Figure 1A). These structures were baited with a two-component aggregation pheromone (3S,6S,7R,10S)-10,11-epoxy-1-bisabolen-3-ol and (3R,6S,7R,10S)-10,11-epoxy-1-bisabolen-3-ol, and pheromone synergist, methyl (2E,4E,6Z)-decatienoate (BMSB High Load Lure®, Trécé Inc.). Lures were changed every 3 months.

The experiment was run as a randomized complete block design with four blocks each one including a pair of similar sites (5-20 ha each). In one of the sites two AK stations ha<sup>-1</sup> were installed, whereas no stations were placed in the control sites. A total of approximately 100 AK stations were set up in the four sites. To evaluate the abundance of BMSB, five black standing pyramid traps (DeadInn®, AgBio Inc.) were set up both in control and AK sites (Figure 1B); the traps were baited with a standard BMSB Monitoring Lure® (Trécé Inc.). Monitoring traps were served weekly from early April to the end of October to count and remove captures. Injuries due to BMSB feeding activity were also scored at the harvest of pear orchards, which were grown in both AK and control sites of three out of four blocks. The other block had peaches and nectarines in both AK and control sites.

Large fluctuations in the total monthly BMSB captures (pooling nymphs and adults) were detected among sites, as shown by the width of the standard errors (Figure 2). The trend of captures was dependent on the season. In spring and early summer no significant differences in BMSB abundance could be detected between AK and



control sites (Figure 2A). On the other hand, from August to October 2021 lower abundances of BMSB were found in the sites with AK station than in control sites (Figure 2B). The decreased BMSB abundance did not turn in a reduction of the injuries to pear as well as to peaches and nectarines.

While the lures were highly effective, some technical issues were reported by the operators who served the AK stations. Recovery capacity of BMSB adults and the needs of prolonged contact of insects with the LLIN seemed crucial as well as quite problematic. Moreover, the active substance alpha-cypermethrin was banned from EU market in 2021. For these reasons, different methods to kill the insects have to be considered. Mass-trapping systems for BMSB relying on glue panels in combination with water traps seem promising and are now under evaluation.




Figure 1. One of the Trinet® pyramid-like structures (A), which were selected as AK stations and the experimental design carried out to test the effects of AK stations on BMSB abundance (B).




Figure 2. Trends of total captures by monitoring traps in the first (A) and in the second (B) half of the sampling season. In the second part of the season a GLMM (with negative binomial error distribution, log link function, months as repeated measures, AR1 covariance type and degrees of freedom corrected by Kenward-Roger method) detected a significant reduction in BMSB abundance ( $p=0.021$ ) in sites where AK stations were deployed (dotted lines) in comparison with control sites (solid lines).

**Key words:** *Halyomorpha halys*, integrated pest management, pheromones, mass-trapping.

**Acknowledgement:** This study was funded by Emilia-Romagna region within the Rural Development Plan 2014-2020 Op. 16.1.01 – GO EIP-Agri - FA 4B, Pr. 'A&K' and coordinated by RI.NOVA.